

Chandra-LETGS observation of Mrk 279: modeling the AGN internal structure

E. Costantini^{*,†}, J.S. Kaastra^{*}, K.C. Steenbrugge^{*}, N. Arav^{**}, J.R. Gabel^{**}
and G. Kriss[‡]

^{*}*SRON, Sorbonnelaan, 2, 8534CA Utrecht, The Netherlands*

[†]*Astronomical Institute, Utrecht University, P.O. Box 80000, 3508TA Utrecht, The Netherlands*

^{**}*CASA, University of Colorado, 389 UCB, Boulder, CO, USA*

[‡]*STSI, 3700 San Martin Drive, Baltimore, MD 21218, USA*

Abstract. We present the *Chandra*-LETGS analysis of Mrk 279, a bright Seyfert 1 galaxy. The spectrum shows a variety of features arising from different physical environments in the vicinity of the black hole. We detect three absorption components, located at the redshift of the source. One of them is likely to arise from the host galaxy. An additional component due to absorption by a collisionally ionized gas at $z=0$ is also observed. Such an absorption can be produced in the outskirts of our Galaxy or in the local group. The emission spectrum is rich in narrow and broad emission features. We tested the hypothesis that the broad emission lines originate in the Broad Line Region (BLR), as studied in the UV band. We find that at least 2 components with different ionization parameters are necessary to simultaneously interpret the broad emission line measured in the UV (by HST-STIS and FUSE) and in the X-rays.

Keywords: X-ray spectra, active galaxies, photoionization of atoms and ions

PACS: 32.30.Rj, 98.54.Cm, 32.80.Fb

INTRODUCTION

The high spectral resolution now available in the X-ray band led to a deeper understanding of the physical processes occurring in active galaxies. Mrk 279 is a type 1 Seyfert galaxy. For these objects, according to the standard model [e.g. 1], our line of sight should intercept an outflowing, highly ionized gas. It is now recognized that many components, differing in ionization level, column density, and outflowing velocity, are actually constituting this outflow [e.g. 6, 9, 13] and that the UV and the X-ray band share at least some of these components [e.g. 13, 10]. Yet, the geometry, the distance, and therefore the density of the X-ray outflow is not yet clear [e.g. 4, 13, 10]. The observation angle for Seyfert 1 objects allows to investigate the innermost parts of the AGN, from the broad line region [BLR, see e.g. 3], at a distance of few light days from the black hole, down to the accretion disk. The bulk of the X-rays are produced here [e.g. 11]. The BLR is likely to produce X-ray radiation, but up to now this radiation was never convincingly observed in X-ray spectra [13, 12].

RESULTS

We processed the *Chandra*-LETGS data of Mrk 279 following the approach described in [7]. The residuals to a basic continuum model show significant excess emission, especially in the 18-24 Å wavelength range, where the main oxygen line features are (O VII triplet and O VIII Ly α). We interpret these emission features as broad lines, possibly arising from the BLR. The best fit shows that the excess can be modeled by Gaussian profiles for O VII, O VIII, N VII, and C VI. Evidence of weaker broad excesses is found for N VI and C V (Fig. 1) and possibly neon (see next paragraph for the modeling). The inclusion of broad emission lines is required to properly fit the absorbed spectrum. The best fit requires indeed absorption by two components of ionized gas. For the lower ionized component we find: $N_H \sim 1.6 \times 10^{22} \text{cm}^{-2}$, and $\log \xi \sim 0.57$, while for the high ionization component the parameters are: $N_H \sim 4 \times 10^{22} \text{cm}^{-2}$ and $\log \xi \sim 2.5$. The two absorbers seem to belong to systems with different outflow velocity, -220_{-90}^{+50} and $-570_{-70}^{+100} \text{ km s}^{-1}$ for the low and high ionization gas, respectively. Another component, whose main features are O I and O II has been modeled by a collisionally ionized gas at a temperature $t \sim 8 \text{ eV}$. Mrk 279 shows also ionized absorption taking place at $z=0$ [8]. In the LETGS band the deepest features of this collisionally ionized component are O VII and C VI.

The Broad Line Region

Thanks to the simultaneous FUSE and HST observation [2], we could compare the X-ray line luminosity with the ones measured for the UV broad emission lines. We used Cloudy [5] for the modeling. The input ionizing spectral continuum was evaluated directly from the HST, FUSE and *Chandra* data. We chose a grid of physical parameters for the BLR clouds and then chose the best fit to our data through χ^2 minimization. In Fig. 2 we display the intrinsic integrated luminosity measured for the X-ray and UV lines (star points) vs the rest frame wavelength of the lines, together with our best fit. Two ionization components for the emitting medium are certainly needed in order to explain the UV and the lower ionization X-ray lines (labeled as “cold” and “warm” in Fig. 2). This result is based mainly on the more precise UV measurements. The higher ionization X-ray lines, such as C VI, N VII, O VIII, Ne IX, and Ne X may require the contribution of a higher ionization component (“hot” in Fig.2). The ionization parameters of these components are roughly $\log \xi \sim -1.5, 1.6$ and 3.6 for the “cold”, “warm”, and “hot” components, respectively. The distance from the ionizing source of the emitting cloud is around 10^{16-17} cm , and the density ranges between 10^9-12 cm^{-3} . The ionization parameter is given by: $\xi = L/nr^2$. Here only the ionizing luminosity L is directly measured. Density effects on the emission may help in determining the parameters in the nr^2 term. However, as a caveat, the r and n values are still coupled.

REFERENCES

1. R. Antonucci, 1993, ARA&A , 31, 473




FIGURE 1. Best fit model for the LETGS data of Mrk 279. The labels refer to the broad lines added in the model.




FIGURE 2. Cloudy prediction of the ionization state of the BLR emitting clouds (see text).

2. N. Arav, J. S. Kaastra, G. A. Kriss, K. T. Korista, J. Gabel, & D. Proga, 2004, astro-ph/0406200
3. J. A. Baldwin, 1997, ASP Conf. Ser. 113: IAU Colloq. 159: Emission Lines in Active Galaxies: New Methods and Techniques, 80
4. E. Behar et al. 2003, ApJ, 598, 232
5. G. J. Ferland, et al. 1998, PASP, 110, 761
6. J. S. Kaastra et al. 2000, A&A, 354, 84
7. J. S. Kaastra et al. 2002, A&A, 386, 427
8. J. S. Kaastra, N. Arav, E. Costantini et al. 2005, in prep.
9. S. Kaspi et al. 2001, ApJ, 554, 216
10. Y. Krongold, F. Nicastro, N. S. Brickhouse, et al. 2003, ApJ, 597, 832
11. K. Nandra, & K. A. Pounds, 1994, MNRAS, 268, 405
12. P. M. Ogle, K. O. Mason, M. J. Page, et al. 2004, ApJ, 606, 151
13. K. C. Steenbrugge et al., 2004, A&A, accepted.